

February 1, 2008

Path integral treatment of a family of super-integrable systems in n -dimensional Euclidean space

M. T. Chefrour

Département de Physique, Faculté des Sciences,
Université Badji Mokhtar, Annaba, Algérie.

F. Benamira and L. Guechi

Laboratoire de Physique Théorique,
Département de Physique, Faculté des Sciences,
Université Mentouri, Route d'Ain El Bey,
Constantine, Algérie.

S. Mameri

Département de Physique, Faculté des Sciences,
Université Ferhat Abbas, Setif, Algérie.

Abstract

The exact path integration for a family of maximally super-integrable systems generalizing the hydrogen atom in the n -dimensional Euclidean space is presented. The Green's function is calculated in parabolic rotational and spherical coordinate systems. The energy spectrum and the correctly normalized wave functions of the bound states are obtained from the poles of the Green's function and their residues, respectively.

PACS 03.65- Quantum theory ; quantum mechanics.

PACS 03.65. Ca - Formalism.

PACS 03.65. Db - Functional analytical methods.

I. Introduction

During the last 40 years, super-integrable dynamical systems have been the object of a considerable interest for their practical importance in various fields. A general classification of Hamiltonians in two and three dimensions possessing dynamical invariance groups was initiated by Smorodinsky, Winternitz and collaborators¹, continued by Kibler and Winternitz², and revived, in recent years, by Evans³. This classification was established according to the number of degrees of freedom, quadratic integrals of motion in the momenta and coordinate systems in which the potential allows the separation of variables. The Hamiltonian systems with these potentials are called super-integrable. Generally, in n dimensions, a system is called "minimally" super-integrable if it has $(2n - 2)$ constants or integrals of motion (including energy), and it is called "maximally" super-integrable if it has $(2n - 1)$ integrals of motion^{4,5}. A list of potentials corresponding to minimally super-integrable and maximally super-integrable systems with corresponding constants of motion in the classical form and all separating coordinate systems has been established by Evans⁶.

In the last decade, Grosche et al have presented a detailed path integral discussion of the so-called Smorodinsky-Winternitz super-integrable Hamiltonians in the three further spaces of constant curvature: two- and three-dimensional Euclidean space⁷, two- and three-dimensional sphere⁸, two- and three-dimensional hyperboloid^{9,10}. Chetouani et al¹¹, whose work followed closely upon that of Grosche and co-workers studied three classes of two- and three-dimensional Smorodinsky-Winternitz super-integrable Hamiltonian systems in Euclidean space using a variant of the $so(2,1)$ algebraic approach.

Recently, a new family of maximally super-integrable systems generalizing the hydrogen atom has been constructed in n -dimensional Euclidean space by Winternitz and Rodriguez¹² and studied in standard quantum mechanics through resolution of Schrödinger equation in parabolic rotational and spherical coordinates. This new family corresponds to the potentials

$$V(x^{(1)}, x^{(2)}, \dots, x^{(n)}) = -\frac{\gamma}{r} + \sum_{i=1}^{n-1} \frac{\beta_i}{(x^{(i)})^2}, \quad (1)$$

where γ and β_i are positive constants and $r = (x^{(1)2} + x^{(2)2} + \dots x^{(n)2})^{\frac{1}{2}}$.

In this paper, we want to present a path integral treatment of these potentials in parabolic rotational and spherical coordinate systems.

our work can be seen as a generalization of the earlier work of Chetouani and Hammann¹³, in which they established only the n -dimensional Coulomb Green function in polar coordinates by means of the Gegenbauer expansion formula.¹⁴

Our study is organized as follows: in the following section, we introduce n -dimensional parabolic rotational coordinates and formulate the path integral associated with the potential $V(x^{(1)}, x^{(2)}, \dots, x^{(n)})$. An explicit Lagrangian path integral formulation is derived. By means of successive applications of appropriate time transformations we reduce the $\{\theta^{(k)}\}$ and (ξ, η) path integrations to the already known ones. The Green function is evaluated in a spectral decomposition. In section III, we construct the Green function in n -dimensional spherical coordinates. The energy spectrum and the normalized wave functions are found as in the case of parabolic rotational coordinates. The section IV will be a conclusion.

II. Parabolic rotational coordinates in E_n and path integral

In n -dimensional parabolic rotational coordinates

$$\begin{aligned}
x^{(1)} &= \xi\eta \cos \theta^{(1)} \cos \theta^{(2)} \dots \cos \theta^{(n-3)} \cos \theta^{(n-2)}, \\
x^{(2)} &= \xi\eta \cos \theta^{(1)} \cos \theta^{(2)} \dots \cos \theta^{(n-3)} \sin \theta^{(n-2)}, \\
x^{(3)} &= \xi\eta \cos \theta^{(1)} \cos \theta^{(2)} \dots \sin \theta^{(n-3)}, \\
&\vdots \\
&\vdots \\
&\vdots \\
x^{(n-2)} &= \xi\eta \cos \theta^{(1)} \sin \theta^{(2)}, \\
x^{(n-1)} &= \xi\eta \sin \theta^{(1)}, \\
x^{(n)} &= \frac{1}{2}(\xi^2 - \eta^2),
\end{aligned} \tag{2}$$

where $0 \leq \xi, \eta < \infty$, $0 \leq \theta^{(\nu)} < \pi$ ($\nu = 2, \dots, n-2$), $0 \leq \theta^{(1)} < 2\pi$, the classical Lagrangian reads as

$$\begin{aligned}\mathcal{L}_{cl} = & \frac{M}{2} \left[(\xi^2 + \eta^2) (\dot{\xi}^2 + \dot{\eta}^2) + \xi^2 \eta^2 \left((\dot{\theta}^{(1)})^2 + \cos^2 \theta^{(1)} (\dot{\theta}^{(2)})^2 + \dots \right. \right. \\ & \left. \left. + \cos^2 \theta^{(1)} \dots \cos^2 \theta^{(n-3)} (\dot{\theta}^{(n-2)})^2 \right) \right] - V, \end{aligned} \quad (3)$$

where the potential $V(x^{(1)}, x^{(2)}, \dots, x^{(n)})$ can be written in the form

$$\begin{aligned}V = & -\frac{2\gamma}{\xi^2 + \eta^2} + \frac{\hbar^2}{2M\xi^2\eta^2} \left[\frac{p_1(p_1 - 1)}{\cos^2 \theta^{(1)} \dots \cos^2 \theta^{(n-2)}} + \frac{p_2(p_2 - 1)}{\cos^2 \theta^{(1)} \dots \sin^2 \theta^{(n-2)}} + \dots \right. \\ & \left. + \frac{p_{n-2}(p_{n-2} - 1)}{\cos^2 \theta^{(1)} \sin^2 \theta^{(2)}} + \frac{p_{n-1}(p_{n-1} - 1)}{\sin^2 \theta^{(1)}} \right], \end{aligned} \quad (4)$$

with

$$p_i = \frac{1}{2} \pm \sqrt{\frac{1}{4} + \frac{2M}{\hbar^2} \beta_i}, \quad i = 1, \dots, n-1. \quad (5)$$

Hence the metric tensor in parabolic rotational coordinates has the form

$$(g_{ab}) = \text{diag}(g_{\xi\xi}, g_{\eta\eta}, g_{\theta^{(1)}\theta^{(1)}}, \dots, g_{\theta^{(n-2)}\theta^{(n-2)}}), \quad (6)$$

where

$$\begin{aligned}g_{\xi\xi} &= g_{\eta\eta} = \xi^2 + \eta^2, & g_{\theta^{(1)}\theta^{(1)}} &= \xi^2 \eta^2 \\ g_{\theta^{(i)}\theta^{(i)}} &= \xi^2 \eta^2 \prod_{k=1}^{i-1} \cos^2 \theta^{(k)}; & i &= 2, \dots, n-2. \end{aligned} \quad (7)$$

The infinitesimal volume element $d^n x$ transforms as follows:

$$d^n x = \sqrt{g} d\xi d\eta d\theta^{(1)} \dots d\theta^{(n-2)}, \quad (8)$$

where

$$\sqrt{g} = \sqrt{\det(g_{ab})} = (\xi^2 + \eta^2)(\xi\eta)^{n-2} \prod_{k=1}^{n-2} (\cos \theta^{(k)})^{n-2-k}. \quad (9)$$

The inverse of (6) is given by

$$(g^{ab}) = (g_{ab})^{-1} = \text{diag}(g_{\xi\xi}^{-1}, g_{\eta\eta}^{-1}, g_{\theta^{(1)}\theta^{(1)}}^{-1}, \dots, g_{\theta^{(n-2)}\theta^{(n-2)}}^{-1}). \quad (10)$$

The momentum operators are

$$\begin{aligned}
P_\xi &= \frac{\hbar}{i} \left(\frac{\partial}{\partial \xi} + \frac{\Gamma_\xi}{2} \right); & \Gamma_\xi &= \frac{2\xi}{\xi^2 + \eta^2} + \frac{n-2}{\xi}, \\
P_\eta &= \frac{\hbar}{i} \left(\frac{\partial}{\partial \eta} + \frac{\Gamma_\eta}{2} \right); & \Gamma_\eta &= \frac{2\eta}{\xi^2 + \eta^2} + \frac{n-2}{\eta}, \\
P_{\theta^{(k)}} &= \frac{\hbar}{i} \left(\frac{\partial}{\partial \theta^{(k)}} + \frac{\Gamma_{\theta^{(k)}}}{2} \right); & \Gamma_{\theta^{(k)}} &= -(n-2-k) \tan \theta^{(k)}, \quad k = 1, \dots, n-2.
\end{aligned} \tag{11}$$

In order to express the quantum Hamiltonian by position and momentum operators, we can choose the Weyl ordering prescription^{15,16,17}, or the product-ordering prescription¹⁸. Due to the special nature of (g_{ab}) and (g^{ab}) respectively, the quantum correction ΔV is the same for both prescriptions. Consequently, we obtain for the Hamiltonian

$$\begin{aligned}
H &= \frac{1}{2M} \left[\frac{1}{\sqrt{\xi^2 + \eta^2}} (P_\xi^2 + P_\eta^2) \frac{1}{\sqrt{\xi^2 + \eta^2}} + \frac{1}{\xi^2 \eta^2} \left[P_{\theta^{(1)}}^2 + \right. \right. \\
&\quad \left. \frac{1}{\cos^2 \theta^{(1)}} \left[P_{\theta^{(2)}}^2 + \frac{1}{\cos^2 \theta^{(2)}} \left[P_{\theta^{(3)}}^2 + \dots + \frac{1}{\cos^2 \theta^{(n-3)}} P_{\theta^{(n-2)}}^2 \right] \dots \right] \right] \\
&\quad - \frac{2\gamma}{\xi^2 + \eta^2} + \frac{\hbar^2}{2M\xi^2\eta^2} \left[\frac{p_1(p_1-1)}{\cos^2 \theta^{(1)} \dots \cos^2 \theta^{(n-2)}} + \frac{p_2(p_2-1)}{\cos^2 \theta^{(1)} \dots \sin^2 \theta^{(n-2)}} + \right. \\
&\quad \left. \dots + \frac{p_{n-2}(p_{n-2}-1)}{\cos^2 \theta^{(1)} \sin^2 \theta^{(2)}} + \frac{p_{n-1}(p_{n-1}-1)}{\sin^2 \theta^{(1)}} \right] + \Delta V, \tag{12}
\end{aligned}$$

with the quantum potential ΔV given by

$$\begin{aligned}
\Delta V &= \frac{\hbar^2}{8M\xi^2\eta^2} \left[(n-2)(n-4) + (n-3)^2 \tan^2 \theta^{(1)} - \frac{2(n-3)}{\cos^2 \theta^{(1)}} \right. \\
&\quad + \frac{1}{\cos^2 \theta^{(1)}} \left[(n-4)^2 \tan^2 \theta^{(2)} - \frac{2(n-4)}{\cos^2 \theta^{(2)}} \right. \\
&\quad + \frac{1}{\cos^2 \theta^{(2)}} \left[(n-5)^2 \tan^2 \theta^{(3)} - \frac{2(n-5)}{\cos^2 \theta^{(3)}} + \dots + \right. \\
&\quad \left. \left. \frac{1}{\cos^2 \theta^{(n-4)}} \left[\tan^2 \theta^{(n-3)} - \frac{2}{\cos^2 \theta^{(n-3)}} \right] \dots \right] \right] \right]. \tag{13}
\end{aligned}$$

In manner similar to that in Refs.^{17,18}, by direct calculation we get for the Lagrangian path integral in the "product form"-definition

$$\begin{aligned}
K(\vec{r}_f, \vec{r}_i; T) &= \int \sqrt{g} \mathcal{D}\xi(t) \mathcal{D}\eta(t) \mathcal{D}\theta^{(1)}(t) \dots \mathcal{D}\theta^{(n-2)}(t) \exp \left\{ \frac{i}{\hbar} \int_0^T (\mathcal{L}_{cl} - \Delta V) dt \right\} \\
&= \lim_{N \rightarrow \infty} \int \prod_{j=1}^N \left(\frac{M}{2i\pi\hbar\varepsilon} \right)^{\frac{n}{2}} \prod_{j=1}^{N-1} (\xi_j^2 + \eta_j^2) (\xi_j \eta_j)^{n-2} \prod_{k=1}^{n-2} (\cos \theta_j^{(k)})^{n-2-k} \\
&\quad \times d\xi_j d\eta_j d\theta_j^{(1)} \dots d\theta_j^{(n-2)} \exp \left\{ \frac{i}{\hbar} \sum_{j=1}^N S^{pc}(j, j-1) \right\}, \tag{14}
\end{aligned}$$

with the short-time action

$$\begin{aligned}
S^{pc}(j, j-1) &= \frac{M}{2\varepsilon} \left[\widehat{(\xi_j^2 + \eta_j^2)} (\Delta^2 \xi_j + \Delta^2 \eta_j) + \widehat{\xi_j^2 \eta_j^2} (\Delta^2 \theta_j^{(1)} \right. \\
&\quad \left. + \cos^2 \theta_j^{(1)} \Delta^2 \theta_j^{(2)} + \dots + \cos^2 \theta_j^{(1)} \dots \cos^2 \theta_j^{(n-3)} \Delta^2 \theta_j^{(n-2)}) \right] \\
&\quad + \frac{2\varepsilon\gamma}{\xi_j^2 + \eta_j^2} - \frac{\varepsilon\hbar^2}{2M\xi_j^2\eta_j^2} \left[-\frac{(n-2)(n-4)}{4} - \frac{(n-3)^2}{4} \right. \\
&\quad \left. + \frac{p_{n-1}(p_{n-1}-1)}{\sin^2 \theta_j^{(1)}} + \frac{(n-3)(n-5)}{4 \cos^2 \theta_j^{(1)}} \right. \\
&\quad \left. + \frac{1}{\cos^2 \theta_j^{(1)}} \left[-\frac{(n-4)^2}{4} + \frac{(n-4)(n-6)}{4 \cos^2 \theta_j^{(2)}} + \frac{p_{n-2}(p_{n-2}-1)}{\sin^2 \theta_j^{(2)}} \right. \right. \\
&\quad \left. + \frac{1}{\cos^2 \theta_j^{(2)}} \left[-\frac{(n-5)^2}{4} + \frac{(n-5)(n-7)}{4 \cos^2 \theta_j^{(3)}} + \frac{p_{n-3}(p_{n-3}-1)}{\sin^2 \theta_j^{(3)}} + \dots \right. \right. \\
&\quad \left. \left. + \frac{1}{\cos^2 \theta_j^{(n-3)}} \left[\frac{p_2(p_2-1)}{\sin^2 \theta_j^{(n-2)}} + \frac{p_1(p_1-1)}{\cos^2 \theta_j^{(n-2)}} \right] \dots \right] \right] \right]. \tag{15}
\end{aligned}$$

In the above we have used the notations: $\varepsilon = t_j - t_{j-1}$, $T = N\varepsilon = t_f - t_i$, $\vec{r}_f = \vec{r}(t_N)$, $\vec{r}_i = \vec{r}(t_0)$, $\widehat{\xi_j^2 + \eta_j^2} = \sqrt{(\xi_j^2 + \eta_j^2)(\xi_{j-1}^2 + \eta_{j-1}^2)}$; $\widehat{\xi_j^2 \eta_j^2} = \xi_j \eta_j \xi_{j-1} \eta_{j-1}$; $\widehat{\cos^2 \theta_j^{(k)}} = \cos \theta_j^{(k)} \cos \theta_{j-1}^{(k)}$.

First, the $\{\theta\}$ path integrations can be entangled with one another by means of pure and appropriate time transformations^{19,20} $t \rightarrow s$ defined by

$dt = \xi^2 \eta^2 \prod_{k=i}^{n-2} \cos^2 \theta^{(n-2-k)} ds; (i = 1, 2, \dots, n-3)$, and carried out using the Pöschl-Teller path integral solution ($0 < \theta < \frac{\pi}{2}$)^{21,22,23,24,25,26}

$$\begin{aligned} K^{(PT)}(\theta_f, \theta_i; T) &= \int \mathcal{D}\theta(t) \exp \left\{ \frac{i}{\hbar} \int_0^T \left[\frac{M}{2} \dot{\theta}^2 - \frac{\hbar^2}{2M} \left(\frac{\kappa^2 - \frac{1}{4}}{\cos^2 \theta} + \frac{\lambda^2 - \frac{1}{4}}{\sin^2 \theta} \right) \right] dt \right\} \\ &= \sum_{J_n=0}^{\infty} \exp \left\{ -\frac{i\hbar T}{2M} (\kappa + \lambda + 2J_n + 1)^2 \right\} \Psi_{J_n}^{(\lambda, \kappa)*}(\theta_i) \Psi_{J_n}^{(\lambda, \kappa)}(\theta_f), \end{aligned} \quad (16)$$

with the normalized wave functions given by

$$\begin{aligned} \Psi_{J_n}^{(\lambda, \kappa)}(\theta) &= \left[2(\kappa + \lambda + 2J_n + 1) \frac{J_n! \Gamma(\kappa + \lambda + J_n + 1)}{\Gamma(\kappa + J_n + 1) \Gamma(\lambda + J_n + 1)} \right]^{\frac{1}{2}} \\ &\times (\sin \theta)^{\lambda + \frac{1}{2}} (\cos \theta)^{\kappa + \frac{1}{2}} P_{J_n}^{(\lambda, \kappa)}(\cos 2\theta), \end{aligned} \quad (17)$$

where $P_{J_n}^{(\lambda, \kappa)}(\cos 2\theta)$ are Jacobi polynomials (¹⁴, p.1035). After performing the integrations over all the angular variables $\theta_j^{(k)}$ ($k = 1, \dots, n-2$), we obtain for the Feynman propagator (14) the following expansion:

$$K(\vec{r}_f, \vec{r}_i; T) = \sum_{\{J_{n-2}\}=0}^{\infty} \Psi_{\{J_{n-2}\}}^* \left(\left\{ \theta_i^{(n-2)} \right\} \right) \Psi_{\{J_{n-2}\}} \left(\left\{ \theta_f^{(n-2)} \right\} \right) K_{\{J_{n-2}\}}(\xi_f, \eta_f, \xi_i, \eta_i; T), \quad (18)$$

where we have abbreviated for $\{J_{n-2}\}$ and $\{\theta^{(n-2)}\}$ the sets $\{J_1, J_2, \dots, J_{n-2}\}$ and $\{\theta^{(1)}, \theta^{(2)}, \dots, \theta^{(n-2)}\}$, respectively. The angular part of the wave functions is found to be

$$\Psi_{\{J_n\}} \left(\left\{ \theta^{(n-2)} \right\} \right) = \prod_{l=1}^{n-2} \Psi_{J_l} \left(\theta^{(n-l-1)} \right), \quad (19)$$

where

$$\begin{aligned} \Psi_{J_l} \left(\theta^{(n-l-1)} \right) &= \left[2 \frac{(m_{l-1} + p_{l+1} + 2J_l + \frac{1}{2}) J_l! \Gamma(m_{l-1} + p_{l+1} + J_l + \frac{1}{2})}{\Gamma(m_{l-1} + J_l + 1) \Gamma(p_{l+1} + J_l + \frac{1}{2})} \right]^{\frac{1}{2}} \\ &\times (\sin \theta^{(n-l-1)})^{p_{l+1}} (\cos \theta^{(n-l-1)})^{m_{l-1} - \frac{1}{2} + 1} \\ &\times P_{J_l}^{(p_{l+1} - \frac{1}{2}, m_{l-1})}(\cos 2\theta^{(n-l-1)}), \end{aligned} \quad (20)$$

with

$$m_0 = p_1 - \frac{1}{2}, \quad m_l = \sum_{i=1}^{l+1} p_i + 2 \sum_{i=1}^l J_i + \frac{l-1}{2}, \quad 1 \leq l \leq n-2. \quad (21)$$

The kernel $K_{\{J_{n-2}\}}(T)$ is given by

$$\begin{aligned} K_{\{J_{n-2}\}}(\xi_f, \eta_f, \xi_i, \eta_i; T) &= (\xi_f \xi_i \eta_f \eta_i)^{\frac{2-n}{2}} (4r_f r_i)^{-\frac{1}{2}} \lim_{N \rightarrow \infty} \int \prod_{j=1}^N \left[\frac{M \sqrt{4r_j r_{j-1}}}{2i\pi \hbar \varepsilon} \right] \\ &\times \prod_{j=1}^{N-1} d\xi_j d\eta_j \exp \left\{ \frac{i}{\hbar} \sum_{j=1}^N A^{pc}(j, j-1) \right\}, \end{aligned} \quad (22)$$

with the short-time action

$$A^{pc}(j, j-1) = \frac{M}{2\varepsilon} (\tilde{\xi}_j^2 + \tilde{\eta}_j^2) (\Delta^2 \xi_j + \Delta^2 \eta_j) + \frac{2\varepsilon \gamma}{\tilde{\xi}_j^2 + \tilde{\eta}_j^2} - \frac{\varepsilon \hbar^2 (\sigma + \frac{n-3}{2})^2 - \frac{1}{4}}{2M \tilde{\xi}_j^2 \tilde{\eta}_j^2}, \quad (23)$$

where we have set

$$\sigma = \sum_{i=1}^{n-1} p_i + 2 \sum_{i=1}^{n-2} J_i, \quad (24)$$

and $\tilde{u}_j = (u_j + u_{j-1})/2$. This brings us to path integration over the coordinates ξ and η . We obtain by introducing the energy E with the help of the Green's function (Fourier transform of the propagator), performing the time transformation $dt = (\xi^2 + \eta^2)ds$ and applying the path integral solution of the radial harmonic oscillator^{27,28} in the ξ and η variable, respectively:

$$\begin{aligned} G(\vec{r}_f, \vec{r}_i; E) &= \int_0^\infty dT \exp\left(\frac{i}{\hbar} E T\right) K(\vec{r}_f, \vec{r}_i; T) \\ &= \sum_{\{J_{n-2}\}=0}^\infty \Psi_{\{J_{n-2}\}}^* \left(\left\{ \theta_i^{(n-2)} \right\} \right) \Psi_{\{J_{n-2}\}} \left(\left\{ \theta_f^{(n-2)} \right\} \right) (\xi_f \xi_i \eta_f \eta_i)^{\frac{3-n}{2}} \\ &\times \left(\frac{M\omega}{i\hbar} \right)^2 \int_0^\infty dS \frac{e^{2\frac{i}{\hbar} \gamma S}}{\sin^2(\omega S)} \exp \left[\frac{iM\omega}{2\hbar} (\xi_f^2 + \eta_f^2 + \xi_i^2 + \eta_i^2) \cot(\omega S) \right] \\ &\times I_{\sigma + \frac{n-3}{2}} \left(\frac{M\omega \xi_f \xi_i}{i\hbar \sin(\omega S)} \right) I_{\sigma + \frac{n-3}{2}} \left(\frac{M\omega \eta_f \eta_i}{i\hbar \sin(\omega S)} \right) \\ &= i\hbar \sum_{N_1, N_2, \{J_{n-2}\}} \frac{\Psi_{N_1, N_2, \{J_{n-2}\}}^* (\xi_i, \eta_i, \{\theta_i^{(n-2)}\}) \Psi_{N_1, N_2, \{J_{n-2}\}} (\xi_f, \eta_f, \{\theta_f^{(n-2)}\})}{E - E_{N_1, N_2, \{J_{n-2}\}}}, \end{aligned} \quad (25)$$

where the properly normalized wave functions and the energy spectrum are given by

$$\begin{aligned} \Psi_{N_1, N_2, \{J_{n-2}\}}(\xi, \eta, \{\theta^{(n-2)}\}) &= \frac{1}{\sqrt{N}} \left(\frac{N_1! N_2!}{\Gamma(N_1 + \sigma + \frac{n-1}{2}) \Gamma(N_2 + \sigma + \frac{n-1}{2})} \right)^{\frac{1}{2}} \\ &\times \left(-\frac{2ME_N}{\hbar^2} \right)^{\frac{n+2\sigma}{4}} (\xi\eta)^\sigma \exp \left[-\sqrt{-\frac{2ME_N}{\hbar^2}} \frac{\xi^2 + \eta^2}{2} \right] \\ &\times L_{N_1}^{\sigma + \frac{n-3}{2}} \left(\sqrt{-\frac{2ME_N}{\hbar^2}} \xi^2 \right) L_{N_2}^{\sigma + \frac{n-3}{2}} \left(\sqrt{-\frac{2ME_N}{\hbar^2}} \eta^2 \right) \\ &\times \Psi_{\{J_{n-2}\}}(\{\theta^{(n-2)}\}), \end{aligned} \quad (26)$$

$$E_N = E_{N_1, N_2, \{J_{n-2}\}} = -\frac{M\gamma^2}{2\hbar^2 N^2}, \quad (27)$$

with $N = N_1 + N_2 + \sigma + \frac{n-1}{2}$. The energy levels and wave functions have been obtained by means of the Hille-Hardy formula (¹⁴, p.1038):

$$\sum_{n=0}^{\infty} \frac{n!}{\Gamma(n + \alpha + 1)} L_n^\alpha(x) L_n^\alpha(y) z^n = \frac{(xyz)^{-\frac{\alpha}{2}}}{1-z} \exp \left(-z \frac{x+y}{1-z} \right) I_\alpha \left(\frac{2\sqrt{xyz}}{1-z} \right); |z| < 1, \quad (28)$$

where $L_n^\alpha(x)$ are the Laguerre polynomials (¹⁴, p.1037).

III. Spherical coordinates in E_n and path integral

We shall now discuss the path integral for this class of super-integrable systems in spherical coordinates. In n -dimensional Euclidean space E_n , the cartesian coordinates $(x^{(1)}, x^{(2)}, \dots, x^{(n)})$, are related to the spherical coordinates $(r, \theta^{(1)}, \theta^{(2)}, \dots, \theta^{(n-1)})$, by means of the following transformation:

$$\begin{aligned} x^{(1)} &= r \cos \theta^{(1)} \\ x^{(2)} &= r \sin \theta^{(1)} \cos \theta^{(2)} \\ &\vdots \\ &\vdots \end{aligned}$$

$$\begin{aligned} x^{(n-1)} &= r \sin \theta^{(1)} \dots \sin \theta^{(n-2)} \cos \theta^{(n-1)} \\ x^{(n)} &= r \sin \theta^{(1)} \dots \sin \theta^{(n-2)} \sin \theta^{(n-1)}, \end{aligned} \quad (29)$$

with $0 < r < \infty, 0 \leq \theta^{(k)} < \pi, k = 1, 2, \dots, n-2, 0 \leq \theta^{(n-1)} < 2\pi$ and $r^2 = \sum_{k=1}^n \left(x^{(k)}\right)^2$. In these spherical coordinates the classical Lagrangian has the form:

$$\begin{aligned} \mathcal{L}_{cl} = & \frac{M}{2} \left[\dot{r}^2 + r^2 \left((\dot{\theta}^{(1)})^2 + \sin^2 \theta^{(1)} (\dot{\theta}^{(2)})^2 + \dots + \sin^2 \theta^{(1)} \dots \sin^2 \theta^{(n-2)} (\dot{\theta}^{(n-1)})^2 \right) \right] \\ & + \frac{\gamma}{r} - \frac{\hbar^2}{2Mr^2} \left[\frac{p_1(p_1-1)}{\cos^2 \theta^{(1)}} + \frac{1}{\sin^2 \theta^{(1)}} \left[\frac{p_2(p_2-1)}{\cos^2 \theta^{(2)}} + \frac{1}{\sin^2 \theta^{(2)}} \left[\frac{p_3(p_3-1)}{\cos^2 \theta^{(3)}} + \dots \right. \right. \right. \\ & \left. \left. \left. + \frac{p_{n-1}(p_{n-1}-1)}{\sin^2 \theta^{(n-2)} \cos^2 \theta^{(n-1)}} \right] \right] \dots \right], \end{aligned} \quad (30)$$

and the metric tensor is given by

$$(g_{ab}) = \text{diag} \left(1, r^2, g_{\theta^{(1)}\theta^{(1)}}, \dots, g_{\theta^{(n-2)}\theta^{(n-2)}} \right), \quad (31)$$

where

$$g_{\theta^{(i)}\theta^{(i)}} = \prod_{k=1}^i \sin^2 \theta^{(k)}, i = 1, 2, \dots, n-2. \quad (32)$$

Of course

$$(g^{ab}) = (g_{ab})^{-1} = \text{diag} \left(1, \frac{1}{r^2}, g_{\theta^{(1)}\theta^{(1)}}^{-1}, \dots, g_{\theta^{(n-2)}\theta^{(n-2)}}^{-1} \right), \quad (33)$$

and

$$\sqrt{g} = \sqrt{\det(g_{ab})} = r^{n-1} \prod_{k=1}^{n-1} \left(\sin \theta^{(k)} \right)^{n-1-k}. \quad (34)$$

The momentum operators have the form

$$\begin{aligned} P_r &= \frac{\hbar}{i} \left(\frac{\partial}{\partial r} + \frac{\Gamma_r}{2} \right); \Gamma_r = \frac{n-1}{r}, \\ P_{\theta^{(k)}} &= \frac{\hbar}{i} \left(\frac{\partial}{\partial \theta^{(k)}} + \frac{\Gamma_{\theta^{(k)}}}{2} \right); \Gamma_{\theta^{(k)}} = (n-1-k) \cot \theta^{(k)}, k = 1, 2, \dots, n-1. \end{aligned} \quad (35)$$

This gives for the Hamiltonian

$$\begin{aligned}
H = & \frac{1}{2M} \left[P_r^2 + \frac{1}{r^2} \left[P_{\theta^{(1)}}^2 + \frac{1}{\sin^2 \theta^{(1)}} \left[P_{\theta^{(2)}}^2 + \frac{1}{\sin^2 \theta^{(2)}} \left[P_{\theta^{(3)}}^2 + \dots + \frac{1}{\sin^2 \theta^{(n-2)}} P_{\theta^{(n-1)}}^2 \right] \dots \right] \right] \right] \\
& - \frac{\gamma}{r} + \frac{\hbar^2}{2Mr^2} \left[\frac{p_1(p_1-1)}{\cos^2 \theta^{(1)}} + \frac{1}{\sin^2 \theta^{(1)}} \left[\frac{p_2(p_2-1)}{\cos^2 \theta^{(2)}} + \frac{1}{\sin^2 \theta^{(2)}} \left[\frac{p_3(p_3-1)}{\cos^2 \theta^{(3)}} + \dots \right. \right. \right. \\
& \left. \left. \left. + \frac{1}{\sin^2 \theta^{(n-2)}} \frac{p_{n-1}(p_{n-1}-1)}{\cos^2 \theta^{(n-1)}} \right] \right] \dots \right] + \Delta V. \tag{36}
\end{aligned}$$

Here the potential ΔV is given by

$$\begin{aligned}
\Delta V = & \frac{\hbar^2}{8Mr^2} \left[(n-1)(n-3) + \frac{1}{\sin^2 \theta^{(1)}} \left[(n-2)^2 \cos^2 \theta^{(1)} - 2(n-2) \right. \right. \\
& \left. \left. + \frac{1}{\sin^2 \theta^{(2)}} \left[(n-3)^2 \cos^2 \theta^{(2)} - 2(n-3) + \frac{1}{\sin^2 \theta^{(3)}} \left[(n-4)^2 \cos^2 \theta^{(3)} \right. \right. \right. \right. \\
& \left. \left. \left. - 2(n-4) + \dots + \frac{1}{\sin^2 \theta^{(n-2)}} \left[\cos^2 \theta^{(n-2)} - 2 \right] \dots \right] \right] \right]. \tag{37}
\end{aligned}$$

Constructing the path integral in E_n , we follow the prescription adopted in the section II and obtain

$$\begin{aligned}
K(\vec{r}_f, \vec{r}_i; T) = & \lim_{N \rightarrow \infty} \int \prod_{j=1}^N \left(\frac{M}{2i\pi\hbar\varepsilon} \right)^{\frac{n}{2}} \prod_{j=1}^{N-1} r_j^{n-1} dr_j \prod_{k=1}^{n-1} \left(\sin \theta_j^{(k)} \right)^{n-1-k} d\theta_j^{(k)} \\
& \times \exp \left\{ \frac{i}{\hbar} \sum_{j=1}^N S^{sc}(j, j-1) \right\}, \tag{38}
\end{aligned}$$

where

$$\begin{aligned}
S^{sc}(j, j-1) = & \frac{M}{2\varepsilon} \left[\Delta^2 r_j + \widehat{r}_j^2 \Delta^2 \theta_j^{(1)} + \widehat{r}_j^2 \widehat{\sin^2 \theta_j^{(1)}} \Delta^2 \theta_j^{(2)} \right. \\
& \left. + \widehat{r}_j^2 \widehat{\sin^2 \theta_j^{(1)}} \dots \widehat{\sin^2 \theta_j^{(n-2)}} \Delta^2 \theta_j^{(n-1)} \right] + \frac{\varepsilon\gamma}{r_j} \\
& - \frac{\varepsilon\hbar^2}{2Mr_j^2} \left[\frac{1}{4} (n-1)(n-3) + \frac{p_1(p_1-1)}{\cos^2 \theta_j^{(1)}} \right. \\
& \left. + \frac{1}{\sin^2 \theta_j^{(1)}} \left[\frac{p_2(p_2-1)}{\cos^2 \theta_j^{(2)}} + \frac{(n-2)^2 \cos^2 \theta_j^{(1)} - 2(n-2)}{4} \right] \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\sin^2 \theta_j^{(2)}} \left[\frac{p_3(p_3 - 1)}{\cos^2 \theta_j^{(3)}} + \frac{(n-3)^2 \cos^2 \theta_j^{(2)} - 2(n-3)}{4} \right. \\
& + \dots + \frac{1}{\sin^2 \theta_j^{(n-2)}} \left[\frac{p_{n-1}(p_{n-1} - 1)}{\cos^2 \theta_j^{(n-1)}} + \frac{\cos^2 \theta_j^{(n-2)} - 2}{4} \right] \dots \left. \right] \Bigg]. \quad (39)
\end{aligned}$$

Path integration in spherical coordinates is quite similar to that carried out in parabolic rotational coordinates. By performing successive time transformations $dt = r^2 \prod_{k=i}^{n-1} \sin^2 \theta^{(n-1-k)} ds$; ($i = 1, 2, \dots, n-2$), and their inverses, we obtain in the separation steps for the angular variables:

$$K(\vec{r}_f, \vec{r}_i; T) = \sum_{\{J_{n-1}\}=0}^{\infty} \Phi_{\{J_{n-1}\}}^* \left(\left\{ \theta_i^{(n-1)} \right\} \right) \Phi_{\{J_{n-1}\}} \left(\left\{ \theta_f^{(n-1)} \right\} \right) K_{\{J_{n-1}\}}(r_f, r_i; T), \quad (40)$$

where $\{J_{n-1}\}$ and $\{\theta^{(n-1)}\}$ denote the sets $\{J_1, J_2, \dots, J_{n-1}\}$ and $\{\theta^{(1)}, \theta^{(2)}, \dots, \theta^{(n-1)}\}$, respectively. The angular part of the wave functions is given by

$$\Phi_{\{J_{n-1}\}} \left(\left\{ \theta^{(n-1)} \right\} \right) = \prod_{k=1}^{n-1} \Phi_{J_k}(\theta^{(k)}), \quad (41)$$

where

$$\begin{aligned}
\Phi_{J_k}(\theta^{(k)}) &= \left[\frac{2(m_{k+1} + p_k + 2J_k + \frac{1}{2}) J_k! \Gamma(m_{k+1} + p_k + J_k + \frac{1}{2})}{\Gamma(p_k + J_k + \frac{1}{2}) \Gamma(m_{k+1} + J_k + 1)} \right]^{\frac{1}{2}} \\
&\times (\sin \theta^{(k)})^{m_{k+1}+1-\frac{n-k}{2}} (\cos \theta^{(k)})^{p_k} P_{J_k}^{(m_{k+1}, p_k - \frac{1}{2})}(\cos 2\theta^{(k)}), \quad (42)
\end{aligned}$$

with

$$m_n = -\frac{1}{2}; \quad m_k = \sum_{i=k}^{n-1} p_i + 2 \sum_{i=k}^{n-1} J_i + \frac{n-k-1}{2}; \quad k = 1, 2, \dots, n-1. \quad (43)$$

The radial path integral $K_{\{J_{n-1}\}}(r_f, r_i; T)$ reads as

$$K_{\{J_{n-1}\}}(r_f, r_i; T) = \frac{1}{(r_f r_i)^{\frac{n-1}{2}}} \int \mathcal{D}r(t) \exp \left\{ \frac{i}{\hbar} \int_0^T \left[\frac{M \dot{r}^2}{2} - \frac{\hbar^2}{2Mr^2} \left(m_1^2 - \frac{1}{4} \right) + \frac{\gamma}{r} \right] dt \right\}. \quad (44)$$

By using the path integral solution for the radial part of the problem of the Coulomb potential^{13,29,30}, the radial Green function is evaluated to be

$$\begin{aligned} G_{\{J_{n-1}\}}(r_f, r_i; E) &= \int_0^\infty dT \exp \left(\frac{iE}{\hbar} T \right) K_{\{J_{n-1}\}}(r_f, r_i; T) \\ &= \frac{M\omega}{i\hbar(r_f r_i)^{\frac{n-2}{2}}} \int_0^\infty dS \frac{e^{\frac{4i\gamma}{\hbar} S}}{\sin(\omega S)} \exp \left\{ \frac{iM\omega}{2\hbar} (r_f + r_i) \cot(\omega S) \right\} \\ &\quad \times I_{2m_1} \left(\frac{M\omega \sqrt{r_f r_i}}{i\hbar \sin(\omega S)} \right) \end{aligned} \quad (45)$$

$$= \frac{1}{i\omega(r_f r_i)^{\frac{n-1}{2}}} \frac{\Gamma(\kappa + m_1 + \frac{1}{2})}{\Gamma(2m_1 + 1)} W_{-\kappa, m_1} \left(\frac{M\omega}{\hbar} r_f \right) M_{-\kappa, m_1} \left(\frac{M\omega}{\hbar} r_i \right), \quad (46)$$

where $\kappa = -2\gamma/\hbar\omega$, $\omega = 2(-2E/M)^{1/2}$ and $r_f > r_i$. The $M_{-\kappa, m_1}(x)$ and $W_{-\kappa, m_1}(x)$ are the Whittaker functions (¹⁴, p. 1059).

Expanding the radial Green function by means of the Hille-Hardy formula (28), we obtain after performing the integration over S in Eq.(45) the energy spectrum and the bound state wave functions, respectively

$$E_N = E_{N_r, \{J_{n-1}\}} = -\frac{M\gamma^2}{2\hbar^2 N^2}; \quad N = N_r + m_1 + \frac{1}{2}, \quad (47)$$

$$\begin{aligned} \Psi_{N_r, \{J_{n-1}\}}(r, \{\theta^{(n-1)}\}) &= \left[\frac{a N_r!}{4\Gamma(N_r + 2m_1 + 1)} \right]^{\frac{1}{2}} \left(\frac{2}{a(N_r + m_1 + \frac{1}{2})} \right)^{m_1 + \frac{3}{2}} r^{m_1 - \frac{n-2}{2}} \\ &\quad \exp \left(-\frac{r}{a(N_r + m_1 + \frac{1}{2})} \right) L_{N_r}^{2m_1} \left(\frac{2r}{a(N_r + m_1 + \frac{1}{2})} \right) \\ &\quad \times \Phi_{\{J_{n-1}\}}(\{\theta^{(n-1)}\}), \end{aligned} \quad (48)$$

with $a = \hbar^2/M\gamma$, and N_r is the radial quantum number.

When we put $\beta_i = 0$ in the expression of $V(x^{(1)}, x^{(2)}, \dots, x^{(n)})$, we obtain the Coulomb potential in n -dimensional Euclidean space. In this case, our results are reduced to those obtained by the path integral approach¹³, or through the resolution of the Schrödinger equation³¹.

IV. Conclusion

In the present work we have shown that the Green function associated to a family of super-integrable systems containing the hydrogen atom as a special case can be calculated by path integral approach in n -dimensional Euclidean space. The explicit path integration has been done in two coordinates systems, namely in parabolic rotational and in spherical coordinates. The $\{\theta^{(k)}\}$ -dependent path integration was manageable by time transformations yielding path integrals of Pöschl-Teller potential forms. It is interesting to note that the construction of the Green function as a spectral expansion gives simultaneously the normalized wave functions and the energy spectrum.

References

- ¹J. Fris, V.Mandrosov, Ya A. Smorodinsky, M.Uhlir and P. Winternitz, *Phys. Lett.* **16** (1965) 354; P.Winternitz, Ya A. Smorodinskii , M. Uhlir and J. Fris, *Yad. Fiz.* **4** (1966) 625; A. A. Makarov , J. A. Smorodinsky , Kh. Valiev and P. Winternitz, *Nuovo Cimento Ser A* **52** (1967) 1061.
- ²M. Kibler and P. Winternitz, *J. Phys. A: Math. Gen.* **20** (1987) 4097; *Phys. Lett. A* **147** (1990) 338.
- ³N. W. Evans, *Phys. Lett. A* **147**(1990) 483; *J. Math. Phys.* **31** (1990) 600; *ibid* **42** (1991) 3369.
- ⁴J. Hietarinta, *Phys. Rep.* **147** (1987) 87.
- ⁵D. Bonatos, C. Daskaloyannis and K. Kokkotas, *Phys. Rev.* **48** (1993) R3407.
- ⁶N. W. Evans, *Phys. Rev. A* **41** (1990) 5666.
- ⁷C. Grosche, G. S. Pogosyan and A. N. Sissakian, *Fortschr. Phys.* **43** (1995) 453.
- ⁸C. Grosche, G. S. Pogosyan and A. N. Sissakian, *Fortschr. Phys.* **43** (1995) 523.
- ⁹C. Grosche, G. S. Pogosyan and A. N. Sissakian, *Phys. Part. Nucl.* **27** (1996) 244.
- ¹⁰C. Grosche, G. S. Pogosyan and A. N. Sissakian, *Phys. Part. Nucl.* **28** (1996) 244.
- ¹¹L. Chetouani, L. Guechi and T. F. Hammann, *J. Math. Phys.* **42** (2001) 4684.
- ¹²P. Winternitz and M. A. Rodriguez, *J. Math. Phys.* **43** (2002) 1309.
- ¹³L. Chetouani and T. F. Hammann, *J. Math. Phys.* **27** (1986) 2944.
- ¹⁴I. S. Gradshteyn and I. M Ryzhik, *Tables of Integrals, Series and Products* (New York: Academic Press, 1965).

- ¹⁵M. Mizrahi, *J. Math. Phys* **16** (1975) 2201.
- ¹⁶T. D. Lee, *Particle Physics and Introduction to Field Theory* (Harwood: Academic Press 1981).
- ¹⁷C. Grosche and F. Steiner, *Zeitschr. Phys. C* **36** (1987)699.
- ¹⁸C. Grosche, *Phys. Lett. A* **128** (1988) 113.
- ¹⁹I. H. Duru and H. Kleinert, *Phys. Lett.B* **84** (1979) 185; *Fortschr. Phys.* **30** (1982) 401.
- ²⁰A. Inomata, *Phys. Lett. A* **87** (1982) 387.
- ²¹M. Böhm and G. Junker, *J. Math. Phys* **28** (1987) 1978.
- ²²I. H. Duru, *Phys Rev. D* **30** (1984) 2121.
- ²³C. Grosche and F. Steiner, *J. Math. Phys* **36** (1995) 2354.
- ²⁴W. Fischer, H. Leschke and P. Müller, *Ann. Phys. (N. Y)* **227** (1993) 206.
- ²⁵A. Inomata, H. Kuratsuji and C. C. Gerry, *Path Integrals and Coherent States of $SU(2)$ and $SU(1,1)$* (Singapore: World Scientific, 1992).
- ²⁶H. Kleinert and I. Mustapic, *J. Math. Phys* **33** (1992) 643.
- ²⁷D. Peak and A. Inomata, *J. Math. Phys* **10** (1969) 1422.
- ²⁸L. Chetouani, L. Guechi and T. F. Hammann, *Phys. Lett. A* **125** (1987) 277.
- ²⁹F. Steiner, *Phys. Lett. A* **106** (1984) 363.
- ³⁰C. Grosche, *Fortschr. Phys.* **40** (1992) 695.
- ³¹L. C. Hostler, *J.Math. Phys.* **11** (1970) 2966.